

OU-HET 410
hep-th/0205036
May 2002

Wilsonian Renormalization Group Approach to $\mathcal{N} = 2$ Supersymmetric Sigma Models

Kiyoshi Higashijima* and Etsuko Itou†

*Department of Physics, Graduate School of Science, Osaka University,
Toyonaka, Osaka 560-0043, Japan*

Abstract

We derive the Wilsonian renormalization group equation in two dimensional $\mathcal{N} = 2$ supersymmetric nonlinear sigma models. This equation shows that the sigma models on compact Einstein Kähler manifolds are asymptotically free. This result is general and does not depend on the specific forms of the Kähler potentials. We also examine the renormalization group flow in a new model which connects two manifolds with different global symmetries.

* E-mail: higashij@phys.sci.osaka-u.ac.jp

† E-mail: itou@het.phys.sci.osaka-u.ac.jp

1 Introduction

Possible interactions in field theories are tightly restricted by the requirement of renormalizability. In four-dimensional scalar field theories, for example, only ϕ^3 and ϕ^4 interactions are renormalizable at least in perturbation theories. In two dimensions, however, nonpolynomial interactions like $\sin \phi$ or four fermion interactions are also renormalizable, and various interesting phenomena such as boson-fermion correspondence or dynamical mass generation are studied without ambiguities due to renormalization. Among field theories with nonpolynomial interactions, nonlinear sigma models provide particularly interesting models in which field variables take values on curved manifolds (target space) like the sphere. Known examples on compact manifolds such as S^N , CP^N or Q^N are asymptotically free and generate the dynamical mass. Some of these theories curiously produce the diquark scalars, gauge fields and fermions as bound states [1].

Recently, the supersymmetric nonlinear sigma models (SNL σ M) on Ricci-flat Kähler manifolds are explicitly constructed [2]. These theories have $\mathcal{N} = 2$ supersymmetry, and are supposed to describe conformally invariant field theories. Although these field theories are interesting for their own sakes, they are also important as possible candidates for the sigma models description of the superstrings propagating in the curved background. Therefore it is curious to study the renormalization properties of $\mathcal{N} = 2$ SNL σ M. In perturbation theories, β function is proportional to the Ricci tensor of the target space in the one loop order [3].

Field theories have rich structure in two dimensions since any polynomial interactions are renormalizable in perturbation theories. Relations among these infinite number of field theories have been studied by using the so-called exact renormalization group equation, which is the Wilsonian renormalization group (WRG) method under the continuous change of the cutoff [4] [5]. In practice, we have to introduce some kind of truncation of the functional differential equation describing the flow of an infinite number of coupling constants. The relevant truncation in the infrared region relies on the derivative expansion of the effective action [6]. The simplest is the local potential approximation, in which one maintains only the potential term without any derivative [7].

In this paper, we derive the Wilsonian renormalization group equation for the in the first nontrivial order of the derivative expansion in $\mathcal{N} = 2$ SNL σ M. Because of the reparametrization invariance of the target manifold, the Lagrangian of the nonlinear sigma model is proportional of the metric of the target manifold and the potential term is absent. Therefore, the first nontrivial order of the derivative expansion is already of the second order in the derivatives [8] [9].

In this order, we obtain the β function for the sigma model

$$\beta(g_{i\bar{j}}) = \frac{1}{2\pi} R_{i\bar{j}} + \gamma \left[\varphi^k g_{i\bar{j},k} + \varphi^{*\bar{k}} g_{i\bar{j},\bar{k}} + 2g_{i\bar{j}} \right]. \quad (1.1)$$

This β function has the one loop correction term and the field rescaling term. We also give the β functions in case of D=3 $\mathcal{N} = 2$ and D=4 $\mathcal{N} = 1$ SNL σ M.

Using this β function, we will show how the target spaces are deformed according to nonperturbative renormalization group flow. We will show, in particular, the theories are asymptotically free when the target spaces are generic compact Einstein Kähler manifolds. We will also construct a new model. The model has two parameters, whose renormalization group flow interpolates two Einstein Kähler manifolds with different global symmetry.

This paper is organized as follows. In section 2, we review the WRG for the most generic action and the approximation method based on the derivative expansion. In section 3, we derive the WRG equation for two dimensional SNL σ M. We also give the explicit WRG equation for the SNL σ M in three and four dimensions. In two dimensions, we show the sigma model with compact Einstein Kähler manifolds as target spaces are asymptotically free in section 4. In section 5, we construct a new model. This model has two parameters in the theory, whose WRG flow interpolates two Einstein Kähler manifolds with different global symmetries.

2 Wilsonian Renormalization Group (WRG): Review

Consider a general Euclidean quantum field theory for the fields Ω_i in D dimensions. The Euclidean path integral is

$$Z = \int [D\Omega_i] \exp [-S[\Omega]]. \quad (2.1)$$

Here $S[\Omega]$ is the most generic Euclidean action, which has the form

$$S[\Omega] \equiv \sum_n \frac{1}{n!} \int_{p_1} \cdots \int_{p_n} \hat{\delta}^{(D)}(p_1 + \cdots + p_n) g_{i_1, \dots, i_n}(p_1, \dots, p_n) \Omega_{i_1}(p_1) \cdots \Omega_{i_n}(p_n), \quad (2.2)$$

where g is a coupling constant, and

$$\hat{\delta}^{(D)} \equiv (2\pi)^D \delta^{(D)} \quad (2.3)$$

is the D dimensional delta-function. The Fourier transformation of Ω is defined by

$$\Omega(x) \equiv \int_p \Omega(p) e^{-ipx} \quad (2.4)$$

in which

$$\int_p \equiv \int \frac{d^D p}{(2\pi)^D}. \quad (2.5)$$

The basic idea to define the effective action is as follows. We divide all fields Ω_i into two group: the high frequency modes ($\Omega_{i>}$) and the low frequency modes ($\Omega_{i<}$). After the higher modes are integrated out, the Wilsonian effective action (S_{eff}) is obtained:

$$\begin{aligned} Z &= \int [D\Omega_i] \exp [-S[\Omega_i]] \\ &= \int [D\Omega_{i>}] [D\Omega_{i<}] \exp [-S[\Omega_{i<}, \Omega_{i>}]] \\ &\equiv \int [D\Omega_{i<}] \exp [-S_{eff}[\Omega_{i<}]]. \end{aligned} \quad (2.6)$$

Our aim is to find the flow of S_{eff} by continuously changing the cutoff Λ which is the boundary between higher and lower momentum. For that purpose, we change the cutoff Λ infinitesimally to $\Lambda(\delta t) = \Lambda e^{-\delta t}$. Thus in the effective action at Λ , we divide all fields into the higher modes $[\Omega_s]$ and the lower modes again. The fields $[\Omega_s]$ which have nonzero value only for the shell momentum between $\Lambda(\delta t)$ and Λ , are integrated out, and we can obtain the effective action for new cutoff $\Lambda(\delta t)$. Hereafter, we simply write S for S_{eff} , and we assume that Z is cutoff independent.

$$\begin{aligned} Z &= \int [D\Omega]_{\Lambda(\delta t)} [D\Omega_s] \exp [-S[\Omega + \Omega_s; \Lambda]] \\ &= \int [D\Omega]_{\Lambda(\delta t)} [D\Omega_s] \exp \left[- \left(S[\Omega; \Lambda] + \frac{\delta S}{\delta \Omega_i} \Omega_s^i + \frac{1}{2} \Omega_s^i \frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \Omega_s^j + O(\Omega_s^3) \right) \right] \\ &= \int [D\Omega]_{\Lambda(\delta t)} \exp \left[- \left(S[\Omega; \Lambda] + \frac{1}{2} \int_{p'} \text{tr} \ln \left(\frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \right) \right. \right. \\ &\quad \left. \left. - \frac{1}{2} \int_{p'} \int_{q'} \frac{\delta S}{\delta \Omega^i} \left(\frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \right)^{-1} \frac{\delta S}{\delta \Omega^j} + O((\delta t)^2) \right) \right] \end{aligned} \quad (2.7)$$

$$\equiv \int [D\Omega]_{\Lambda(\delta t)} \exp [-S[\Omega; \Lambda(\delta t)]], \quad (2.8)$$

where

$$\int_{p'} = \int \frac{d\Omega_D}{(2\pi)^D} \int_{\Lambda(\delta t)}^{\Lambda} dp' \sim O(\delta t), \quad (2.9)$$

with $\int d\Omega_D$ being the surface integral of the D-dimensional unit sphere. We can drop terms of order $O(\Omega_s^3)$, because such terms vanish when $\delta t \rightarrow 0$. In

Eq.(2.7) the action $S[\Omega; \Lambda]$ is obtained from $S[\Omega; \Lambda]$ by dropping all fields with momentum above $\Lambda(\delta t)$

$$S[\Omega; \Lambda] = \sum_n \frac{1}{n!} \int_{p_1} \cdots \int_{p_n} \hat{\delta}(p_1 + \cdots + p_n) g(\Lambda) \Omega_{i_1} \cdots \Omega_{i_n}. \quad (2.10)$$

On the other hand, the action $S[\Omega; \Lambda(\delta t)]$ in Eq.(2.8) depend on the fields with the lower momentum, and the coupling constants which are defined at the lower cutoff $\Lambda(\delta t)$:

$$S[\Omega; \Lambda(\delta t)] = \sum_n \frac{1}{n!} \int_{p_1} \cdots \int_{p_n} \hat{\delta}(p_1 + \cdots + p_n) g(\Lambda(\delta t)) \Omega_{i_1} \cdots \Omega_{i_n}. \quad (2.11)$$

The difference between Eq.(2.10) and Eq.(2.11) lies in the variation of coupling constants when the cutoff is changed.

$$\begin{aligned} & S[\Omega; \Lambda(\delta t)] - S[\Omega; \Lambda] \\ &= \sum_n \frac{1}{n!} \int_{p_1} \cdots \int_{p_n} \hat{\delta}(p_1 + \cdots + p_n) [g(\Lambda(\delta t)) - g(\Lambda)] \Omega_{i_1} \cdots \Omega_{i_n} \\ &= -\delta t \sum_n \frac{1}{n!} \int_{p_1} \cdots \int_{p_n} \hat{\delta}(p_1 + \cdots + p_n) \left(\Lambda \frac{\partial}{\partial \Lambda} g(\Lambda) \right) \Omega_{i_1}(p_1) \cdots \Omega_{i_n}(p_n) \\ &= \frac{1}{2} \int_{p'} \text{tr} \ln \left(\frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \right) - \frac{1}{2} \int_{p'} \int_{q'} \frac{\delta S}{\delta \Omega^i} \left(\frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \right)^{-1} \frac{\delta S}{\delta \Omega^j}. \end{aligned} \quad (2.12)$$

If the coupling constants depend on momenta, this cutoff dependence of the coupling constant is given by

$$\Lambda \frac{\partial}{\partial \Lambda} g(\Lambda) = \Lambda \frac{d}{d\Lambda} g(\Lambda) - \sum_i p_i^\mu \frac{\partial}{\partial p_i^\mu} g(\Lambda). \quad (2.13)$$

In order to derive a differential equation for S , we transform all fields and coupling constants to dimensionless quantities. The mass dimension of a coupling constant g is given

$$\dim[g] = D - \sum_{\Omega_i} (d_{\Omega_i} + \gamma_{\Omega_i}), \quad (2.14)$$

where D denotes space-time dimensions, and d_{Ω_i} , γ_{Ω_i} are the canonical and anomalous dimensions of Ω_i . Since the cutoff dependence of the coupling constant can be written

$$\Lambda \frac{d}{d\Lambda} g = \Lambda \frac{\partial}{\partial \Lambda} g - \dim[g] \cdot g + \sum_i p_i^\mu \frac{\partial}{\partial p_i^\mu} g(\Lambda), \quad (2.15)$$

the WRG equation for the effective action S is

$$\Lambda \frac{d}{d\Lambda} S = -\frac{1}{\delta t} [S[\Omega; \Lambda(\delta t)] - S[\Omega; \Lambda]]$$

$$\begin{aligned}
& - \left[D - \sum_{\Omega_i} \int_p \hat{\Omega}_i(p) \left(d_{\Omega_i} + \gamma_{\Omega_i} + \hat{p}^\mu \frac{\partial}{\partial \hat{p}^\mu} \right) \frac{\delta}{\delta \hat{\Omega}_i(p)} \right] \hat{S} \\
& \equiv - \frac{d}{dt} \hat{S},
\end{aligned} \tag{2.16}$$

where the caret indicates dimensionless quantities.

Using Eq.(2.12), we obtain WRG equation for dimensionless action:

$$\begin{aligned}
\frac{d}{dt} S[\Omega; t] &= \frac{1}{2\delta t} \int_{p'} \text{tr} \ln \left(\frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \right) \\
& - \frac{1}{2\delta t} \int_{p'} \int_{q'} \frac{\delta S}{\delta \Omega^i(p')} \left(\frac{\delta^2 S}{\delta \Omega^i(p') \delta \Omega^j(q')} \right)^{-1} \frac{\delta S}{\delta \Omega^j(q')} \\
& + \left[D - \sum_{\Omega_i} \int_p \hat{\Omega}_i(p) \left(d_{\Omega_i} + \gamma_{\Omega_i} + \hat{p}^\mu \frac{\partial}{\partial \hat{p}^\mu} \right) \frac{\delta}{\delta \hat{\Omega}_i(p)} \right] \hat{S}.
\end{aligned} \tag{2.17}$$

The terms

$$\int_{p'} \text{tr} \ln \left(\frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \right) \tag{2.18}$$

and

$$\int_{p'} \int_{q'} \frac{\delta S}{\delta \Omega^i(p')} \left(\frac{\delta^2 S}{\delta \Omega^i(p') \delta \Omega^j(q')} \right)^{-1} \frac{\delta S}{\delta \Omega^j(q')} \tag{2.19}$$

correspond to the one-loop and dumbbell diagrams respectively shown in Figures 1 and 2.




Figure 1: one-loop diagram




Figure 2: dumbbell diagram

Here, the single and double lines represent the lower and higher modes, and the latter are integrated out in Eq.(2.17). In the limit $\delta t \rightarrow 0$, the contribution of the higher order diagrams disappears.

If there are fermionic fields in addition to bosonic fields, the WRG equation (2.17) can be rewritten:

$$\begin{aligned} \frac{d}{dt} S[\Omega; t] &= \frac{1}{2\delta t} \int_{p'} str \ln \left(\frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \right) \\ &\quad - \frac{1}{2\delta t} \int_{p'} \int_{q'} (-1)^F \frac{\delta S}{\delta \Omega^i(p')} \left(\frac{\delta^2 S}{\delta \Omega^i(p') \delta \Omega^j(q')} \right)^{-1} \frac{\delta S}{\delta \Omega^j(q')} \\ &\quad + \left[D - \sum_{\Omega^i} \int_p \hat{\Omega}^i(p) (d_{\Omega^i} + \gamma_{\Omega^i} + \hat{p}^\mu \frac{\partial}{\partial \hat{p}^\mu}) \frac{\delta}{\delta \hat{\Omega}^i(p)} \right] \hat{S}, \end{aligned} \quad (2.20)$$

where

$$\begin{aligned} str \ln \left(\frac{\delta^2 S}{\delta \Omega^i \delta \Omega^j} \right) &= str \ln M_{i\bar{j}} \\ &= str \ln \begin{pmatrix} M_{BB} & M_{BF} \\ M_{FB} & M_{FF} \end{pmatrix} \\ &= tr \ln M_{BB} - tr \ln N_{FF}, \end{aligned} \quad (2.21)$$

with

$$N_{FF} = M_{FF} - M_{FB} M_{BB}^{-1} M_{BF}. \quad (2.22)$$

The action $S[\Omega; t]$ in Eq.(2.17) includes an infinite number of coupling constants $g_{i_1, \dots, i_n}$, and Eq.(2.17) gives infinite number of differential equations among them. To make these equations more tractable, we usually expand the effective action in powers of derivatives and maintain the first few terms. We often introduce a symmetry (supersymmetry, gauge symmetry, Z_2 symmetry etc) to further decrease the number of independent coupling constants.

Consider, for example, a single real scalar field theory which is invariant under $\varphi \rightarrow -\varphi$ (Z_2 symmetry). We can expand the most generic effective

action as:

$$S[\varphi] = \int d^D x \left[V[\varphi] + \frac{1}{2} K[\varphi] (\partial_\mu \varphi)^2 + H_1[\varphi] (\partial_\mu \varphi)^4 + H_2[\varphi] (\partial_\mu \partial^\mu \varphi)^2 + \dots \right],$$

where $V[\varphi], K[\varphi], H_1[\varphi], \dots$ are functions of φ . To second order of derivatives, this action becomes

$$S[\varphi] = \int d^D x \left[V[\varphi] + \frac{1}{2} K[\varphi] (\partial_\mu \varphi)^2 \right]. \quad (2.23)$$

Substituting this $S[\varphi]$ into Eq.(2.17) and expanding the right hand side of Eq.(2.17) up to $O(\partial^2)$, we obtain two differential equations for $V[\varphi], K[\varphi]$.

In scalar field theories, one often maintains only the local potential term $V[\varphi]$. In this paper, we consider $\mathcal{N} = 2$ SNL σ M and work at the second order of derivatives for the scalar fields. In these theories, the local potential term is forbidden by the reparametrization invariance of the target manifold.

3 Supersymmetric Nonlinear Sigma Model

For $\mathcal{N} = 2$ supersymmetric nonlinear sigma model (SNL σ M) in two dimensions (D=2), the action is determined by the Kähler potential $K[\Phi, \Phi^\dagger]$:

$$S = \int dV K[\Phi, \Phi^\dagger], \quad (3.1)$$

where

$$\int dV \equiv \int d^2 x d^2 \theta d^2 \bar{\theta}.$$

This is also true for $\mathcal{N} = 2$ SNL σ M in D=3 and $\mathcal{N} = 1$ SNL σ M in D=4, then we replace the space-time dimensions 2 with D . Here, Φ is a chiral superfield and can be written by component fields:

$$\begin{aligned} \Phi^i(y) &= \varphi^i(y) + \sqrt{2} \theta \psi^i(y) + \theta \theta F^i(y) \\ &= \varphi^i(x) + i \theta \sigma^\mu \bar{\theta} \partial_\mu \varphi^i(x) + \frac{1}{4} \theta \theta \bar{\theta} \bar{\theta} \partial^\mu \partial_\mu \varphi^i(x) \\ &\quad + \sqrt{2} \theta \psi^i(x) - \frac{i}{\sqrt{2}} \theta \theta \partial_\mu \psi^i(x) \sigma^\mu \bar{\theta} + \theta \theta F^i(x) \end{aligned} \quad (3.2)$$

$$\equiv \varphi^i(x) + \delta \Phi^i(x), \quad (3.3)$$

where,

$$y^\mu = x^\mu + i \theta \sigma^\mu \bar{\theta} \quad (3.4)$$

Using (3.3), we expand the action (3.1) around scalar fields (φ, φ^*) . It is sufficient to expand it to $O((\delta\Phi)^4)$, because $\delta\Phi$ (and $\delta\Phi^\dagger$) contains at least one θ . Integrating out over Grassmann numbers (θ and $\bar{\theta}$), we obtain the SNL σ M action written by component fields:

$$S = \int d^D x \left[g_{n\bar{m}} \left(\partial^\mu \varphi^n \partial_\mu \varphi^{*\bar{m}} + \frac{i}{2} \bar{\psi}^{\bar{m}} \sigma^\mu (D_\mu \psi)^n + \frac{i}{2} \psi^n \bar{\sigma}^\mu (D_\mu \bar{\psi})^{\bar{m}} + \bar{F}^{\bar{m}} F^n \right) - \frac{1}{2} K_{,n\bar{m}\bar{l}} \bar{F}^{\bar{l}} \psi^n \psi^{\bar{m}} - \frac{1}{2} K_{,n\bar{m}\bar{l}} F^n \bar{\psi}^{\bar{m}} \bar{\psi}^{\bar{l}} + \frac{1}{4} K_{,nm\bar{k}\bar{l}} (\bar{\psi}^{\bar{k}} \bar{\psi}^{\bar{l}}) (\psi^n \psi^{\bar{m}}) \right]. \quad (3.5)$$

First, we consider the bosonic part of the action (3.5). The loop correction term in the WRG eq.(2.17) cannot be written in covariant form in general. We use the Kähler normal coordinates (KNC) expansion of the action to obtain the covariant expression for the loop correction [10] [11].

The Kähler normal coordinates (ω, ω^*) are defined by the following condition:

$$K_{,\bar{j}i_1 \dots i_n}(\omega, \omega^*)|_0 = 0. \quad (3.6)$$

Here, the index "0" denotes a value evaluated at the origin of KNC, which is the expansion point.

The Riemann normal coordinates (RNC) are well known in perturbative calculations [12] [13]. All geodesics in RNC become straight lines. The RNC in Kähler manifolds, however, is not chiral and the coordinate transformation from the holomorphic coordinates to RNC is not holomorphic. On the contrary, in KNC, geodesics cannot become straight lines, but the KNC preserves holomorphy.

Let us decompose arbitrary holomorphic coordinates (z^i) into a background field φ^i and small fluctuation π^i around it: $z^i = \varphi^i + \pi^i$. The coordinate transformation from coordinates z^i to KNC ω is given by

$$\omega^i = \pi^i + \sum_{n=2}^{\infty} \frac{1}{n!} [g^{i\bar{j}} K_{,\bar{j}i_1 \dots i_n}(z^i, z^{*\bar{i}})]_\varphi \pi^{i_1} \dots \pi^{i_n}. \quad (3.7)$$

These KNC ω^i is known to be transformed as the holomorphic tangent vectors. Since they have the well-defined transformation property, we can use convenient coordinate system for our calculation.

In the D dimensional sigma model action

$$S_{scalar} = \int d^D x g_{i\bar{j}}(\varphi, \varphi^*) \partial^\mu \varphi^i \partial_\mu \varphi^{*\bar{j}}, \quad (3.8)$$

the background fields φ^i are considered as the lower frequency modes and the fluctuation π^i as the higher frequency modes. The Euclidean path integral Z in Eq.(2.7) is written

$$Z = \int [D\varphi][D\varphi^*][D\pi][D\bar{\pi}] \exp \left[-S[\varphi + \pi, \varphi^* + \bar{\pi}] \right]. \quad (3.9)$$

We regard the background field φ^i as the origin of KNC, and transform the fluctuations π^i to KNC fields $\bar{\pi}^i$:

$$Z \xrightarrow{KNC} \int [D\varphi][D\varphi^*][D\bar{\pi}][D\bar{\pi}^*] \exp \left[-\bar{S}[\varphi + \bar{\pi}, \varphi^* + \bar{\pi}^*] \right], \quad (3.10)$$

where bars indicate quantities in KNC, and $\bar{S}$ is given by

$$\bar{S}[\varphi + \bar{\pi}, \varphi^* + \bar{\pi}^*] = \int d^D x \bar{g}_{i\bar{j}}(\varphi + \bar{\pi}, \varphi^* + \bar{\pi}^*) \partial_\mu (\varphi + \bar{\pi})^i \partial^\mu (\varphi^* + \bar{\pi}^*)^{\bar{j}} \quad (3.11)$$

The KNC fields $\bar{\pi}(x)$ can be expanded by tangent vectors $\bar{\omega}(x)$, by solving Eq.(3.7) for π^i and taking KNC system

$$\bar{\pi}(x) = \bar{\omega}^i(x) - \frac{1}{2} \bar{\Gamma}_{k_1 k_2}^i|_\varphi \bar{\omega}^{k_1}(x) \bar{\omega}^{k_2}(x) + O(\bar{\omega}^3). \quad (3.12)$$

Substituting Eq.(3.12) into Eq.(3.11) and using the property of KNC (3.6), we obtain the expansion of the sigma model action to the second order of the fluctuations in KNC:

$$\begin{aligned} \bar{S}[\varphi + \bar{\pi}, \varphi^* + \bar{\pi}^*] = & \int d^D x \bar{g}_{i\bar{j}}|_\varphi \left(\partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}} + \partial_\mu \varphi^i \partial^\mu \bar{\omega}^{*\bar{j}} + \partial_\mu \bar{\omega}^i \partial^\mu \varphi^{*\bar{j}} + \partial_\mu \bar{\omega}^i \partial^\mu \bar{\omega}^{*\bar{j}} \right) \\ & + \bar{g}_{i\bar{j}, k\bar{l}}|_\varphi \left(\partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}} \bar{\omega}^k \bar{\omega}^{*\bar{l}} + \frac{1}{2} \partial_\mu \varphi^i \partial^\mu \varphi^k \bar{\omega}^{*\bar{j}} \bar{\omega}^{*\bar{l}} + \frac{1}{2} \partial_\mu \varphi^{*\bar{j}} \partial^\mu \varphi^{*\bar{l}} \bar{\omega}^i \bar{\omega}^k \right) \\ & + O(\bar{\omega}^3). \end{aligned} \quad (3.13)$$

From this expansion, we can read the matrix

$$\left(\frac{\delta^2 S}{\delta \Omega^i(p') \delta \Omega^j(q')} \right) = M_{ij}(q', p') \quad (3.14)$$

in Eq.(2.17), whose trace corresponds to the one loop correction term. These matrix elements are in momentum space

$$M(q', p') = \begin{pmatrix} M_{k\bar{l}} & M_{\bar{k}\bar{l}} \\ M_{kl} & M_{\bar{k}l} \end{pmatrix}, \quad (3.15)$$

where

$$\begin{aligned} M_{k\bar{l}} & \equiv \frac{\delta^2 \bar{S}}{\delta \varphi^{*\bar{l}}(q') \delta \varphi^k(p')} \\ & = \int_{p', q', P} (-p' \cdot q') \bar{g}_{k\bar{l}}(P) \hat{\delta}(p' + q' + P) \\ & \quad + \int_{p', q', p, q, P''} (-p \cdot q) \bar{g}_{i\bar{j}, k\bar{l}}(P'' + p' + q') \varphi^i(p) \varphi^{*\bar{j}} \hat{\delta}(p' + q' + p + q + P''), \end{aligned} \quad (3.16)$$

$$M_{\bar{k}\bar{l}} = \int_{p',q',p,q,P''} (-p \cdot q) \bar{g}_{i\bar{k},j\bar{l}}(P'' + p' + q') \varphi^i(p) \varphi^j(q) \hat{\delta}(p' + q' + p + q + P''), \quad (3.17)$$

$$M_{kl} = \int_{p',q',p,q,P''} (-p \cdot q) \bar{g}_{k\bar{i},l\bar{j}}(P'' + p' + q') \varphi^{*\bar{i}}(p) \varphi^{*\bar{j}}(q) \hat{\delta}(p' + q' + p + q + P''), \quad (3.18)$$

$$M_{\bar{k}l} = \int_{p',q',P} (-p' \cdot q') \bar{g}_{l\bar{k}}(P) \hat{\delta}(p' + q' + P) + \int_{p',q',p,q,P''} (-p \cdot q) \bar{g}_{i\bar{j},l\bar{k}}(P'' + p' + q') \varphi^i(p) \varphi^{*\bar{j}} \hat{\delta}(p' + q' + p + q + P''). \quad (3.19)$$

We write the (1,1) element (3.16) as

$$M_{k\bar{l}} \equiv \langle \bar{l}; -q' | \hat{M}_1 | k; p' \rangle, \quad (3.20)$$

to define the operator $\hat{M}_1$. Here, the states $|k; p\rangle$ are defined by

$$\langle k; p | \bar{l}; q \rangle = \langle \bar{l}; p | k; q \rangle = \hat{\delta}(q - p) \delta_{k\bar{l}}, \quad (3.21)$$

$$\sum_p \sum_{k\bar{l}} \left[|k; p\rangle \delta^{\bar{l}k} \langle \bar{l}; p| + |\bar{l}; p\rangle \delta^{\bar{l}k} \langle k; p| \right] = 1. \quad (3.22)$$

From Eq.(3.20), the explicit form of the operator $\hat{M}_1$ is

$$(\hat{M}_1)_{k\bar{l}} = \hat{p} \bar{g}_{k\bar{l}} \hat{p} + \bar{g}_{i\bar{j},k\bar{l}} \partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}}. \quad (3.23)$$

Similarly, from the other elements of the matrix (3.17),(3.18),(3.19) we can define the following operators:

$$(\hat{M}_2)_{\bar{k}\bar{l}} = \bar{g}_{i\bar{k},j\bar{l}} \partial_\mu \varphi^i \partial^\mu \varphi^j, \quad (3.24)$$

$$(\hat{M}_3)_{kl} = \bar{g}_{k\bar{i},l\bar{j}} \partial_\mu \varphi^{*\bar{i}} \partial^\mu \varphi^{*\bar{j}}, \quad (3.25)$$

$$(\hat{M}_4)_{\bar{k}l} = \hat{p} \bar{g}_{\bar{k}l} \hat{p} + \bar{g}_{i\bar{j},k\bar{l}} \partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}}. \quad (3.26)$$

The one loop correction term to $O(\partial^2)$ can be decomposed as follows:

$$\int_{p'} \text{tr} \ln M_{ij}(-p', p') = \int_{p'} \text{tr} \ln \hat{M}_{1i\bar{j}} + \int_{p'} \text{tr} \ln \hat{M}_{4i\bar{j}}, \quad (3.27)$$

because $\hat{M}_2$ and $\hat{M}_3$ contribute only to higher derivative terms. Noting that $\bar{g}_{i\bar{j},k}$ and $\bar{g}_{i\bar{j},\bar{k}}$ are zero in KNC, we can calculate this trace:

$$\begin{aligned} \int_{p'} \text{tr} \ln M_{ij}(-p', p') &= \frac{2}{(2\pi)^D} (\delta t) \int d\Omega_D \int d^D x \text{tr} \ln \bar{g}_{i\bar{j}} \\ &+ \frac{2}{(2\pi)^D} (\delta t) \int d\Omega_D \int d^D x \bar{g}_{i\bar{j},k\bar{l}} \bar{g}^{k\bar{l}} \partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}}. \end{aligned} \quad (3.28)$$

Now we have to discuss the contribution of the fermion part in the supersymmetric action (3.5). It turns out that there is no fermion contribution to the bosonic action except for the first term in (3.28) with the opposite sign. Therefore, the first term in the right hand side of Eq.(3.28) cancels with the similar contribution from the fermionic part. In KNC, the second term can be written as

$$-\frac{2}{(2\pi)^D}(\delta t) \int d\Omega_D \int d^D x \bar{R}_{i\bar{j}} \partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}}. \quad (3.29)$$

Finally, we transform this result from KNC to the original coordinates. Then, the first term of the WRG eq.(2.17) is

$$\frac{1}{2\delta t} \int_{p'} \text{tr} \ln M_{ij} = -\frac{1}{(2\pi)^D} \int d\Omega_D \int d^D x R_{i\bar{j}} \partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}}. \quad (3.30)$$

The second term of Eq.(2.17), the contribution of the dumbbell diagram, vanishes to the order $O(\partial^2)$ in the derivative expansion, because external lines carrying soft momenta in the derivative expansion cannot satisfy the energy-momentum conservation law with the hard shell-momenta carried by the internal double lines in the dumbbell diagram.

Similar derivation can be applied to $D=2$ $\mathcal{N}=2$, $D=3$ $\mathcal{N}=2$ and $D=4$ $\mathcal{N}=1$ SNL σ M. We obtain the WRG equations for scalar part action for these SNL σ M.

- In the case of $D=2$ $\mathcal{N}=2$ SNL σ M

$$\begin{aligned} & \frac{d}{dt} \int d^2 x g_{i\bar{j}} (\partial_\mu \varphi)^i (\partial^\mu \varphi^*)^{\bar{j}} \\ &= \int d^2 x \left[-\frac{1}{2\pi} R_{i\bar{j}} - \gamma \left[\varphi^k g_{i\bar{j},k} + \varphi^{*\bar{k}} g_{i\bar{j},\bar{k}} + 2g_{i\bar{j}} \right] \right] (\partial_\mu \varphi)^i (\partial^\mu \varphi^*)^{\bar{j}}. \end{aligned} \quad (3.31)$$

- In the case of $D=3$ $\mathcal{N}=2$ SNL σ M

$$\begin{aligned} & \frac{d}{dt} \int d^3 x g_{i\bar{j}} (\partial_\mu \varphi)^i (\partial^\mu \varphi^*)^{\bar{j}} \\ &= \int d^3 x \left[-\frac{1}{2\pi^2} R_{i\bar{j}} - \gamma \left[\varphi^k g_{i\bar{j},k} + \varphi^{*\bar{k}} g_{i\bar{j},\bar{k}} + 2g_{i\bar{j}} \right] \right. \\ & \quad \left. - \frac{1}{2} \left[\varphi^k g_{i\bar{j},k} + \varphi^{*\bar{k}} g_{i\bar{j},\bar{k}} \right] \right] (\partial_\mu \varphi)^i (\partial^\mu \varphi^*)^{\bar{j}}. \end{aligned} \quad (3.32)$$

- In the case of D=4 $\mathcal{N} = 1$ SNL σ M

$$\begin{aligned}
& \frac{d}{dt} \int d^4x g_{i\bar{j}} (\partial_\mu \varphi)^i (\partial^\mu \varphi^*)^{\bar{j}} \\
&= \int d^4x \left[-\frac{1}{8\pi^2} R_{i\bar{j}} - \gamma \left[\varphi^k g_{i\bar{j},k} + \varphi^{*\bar{k}} g_{i\bar{j},\bar{k}} + 2g_{i\bar{j}} \right] \right. \\
&\quad \left. - \left[\varphi^k g_{i\bar{j},k} + \varphi^{*\bar{k}} g_{i\bar{j},\bar{k}} \right] \right] (\partial_\mu \varphi)^i (\partial^\mu \varphi^*)^{\bar{j}}. \tag{3.33}
\end{aligned}$$

In two dimensions, for example, the β function of the Kähler metric is

$$\begin{aligned}
\frac{d}{dt} g_{i\bar{j}} &= -\frac{1}{2\pi} R_{i\bar{j}} - \gamma \left[\varphi^k g_{i\bar{j},k} + \varphi^{*\bar{k}} g_{i\bar{j},\bar{k}} + 2g_{i\bar{j}} \right] \\
&\equiv -\beta(g_{i\bar{j}}). \tag{3.34}
\end{aligned}$$

We concentrated to the discussion of the WRG equation for the bosonic part of the action. In supersymmetric theories, Eq.(3.34) has to be the scalar part of the relation among superfields. Other parts will have contributions both from bosonic and fermionic parts. Here, we simply assume supersymmetry to derive supersymmetric relation. In this paper, we used straight cut-off in momentum space to obtain the WRG equation. We will need some modification or counter terms to maintain supersymmetry. If we assume supersymmetry, the WRG equation for Kähler potential should be

$$\begin{aligned}
\frac{d}{dt} \int dV K[\Phi, \Phi^\dagger] &= \int dV \Delta K_1[\Phi, \Phi^\dagger] \\
&+ \left[2 - \sum_{\Omega^i} \int_p \hat{\Omega}^i(p) (d_{\Omega^i} + \gamma_{\Omega^i} + \hat{p}^\mu \frac{\partial}{\partial \hat{p}^\mu}) \frac{\delta}{\delta \hat{\Omega}^i(p)} \right] \hat{S}, \tag{3.35}
\end{aligned}$$

where Ω stands for $\varphi, \varphi^*, \psi, \bar{\psi}, F, \bar{F}$, and ΔK_1 is the one-loop correction:

$$\Delta K_1 = \frac{1}{2\pi} \ln \det g_{k\bar{l}}[\Phi, \Phi^\dagger]. \tag{3.36}$$

Expanding Eq.(3.35) around scalar fields, as in Eq.(3.5), we can obtain the WRG equations for various terms in Eq.(3.5). These equations are consistent with Eq.(3.31).

From now on, we consider only D=2 $\mathcal{N} = 2$ SNL σ M.

4 Generic Einstein Kähler Manifolds

Using the WRG equation for SNL σ M action (3.31), we find that SNL σ M are asymptotically free, when target spaces are the compact Einstein Kähler manifolds.

The Einstein Kähler manifolds satisfy the following condition:

$$R_{i\bar{j}} = \frac{h}{a^2} g_{i\bar{j}}, \quad (4.1)$$

where a is the radius of the manifold, which is related to the coupling constant λ by

$$\lambda \equiv \frac{1}{a}. \quad (4.2)$$

If the manifold is Hermitian symmetric space (G/H), the positive constant h in Eq.(4.1) is the eigenvalue of the quadratic Casimir operator in adjoint representation of global symmetry G [2]. For the $CP^N = SU(N+1)/[SU(N) \times U(1)]$ model, its value is $N+1$. We will discuss this model later.

When the manifolds have the radius $a = \frac{1}{\lambda}$, the scalar part of SNL σ M Lagrangian can be represented in the following form:

$$\begin{aligned} \mathcal{L}_{scalar} &= g_{i\bar{j}}(\varphi, \varphi^*) \partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}} \\ &\xrightarrow{\varphi, \varphi^* \approx 0} \frac{1}{\lambda^2} \delta_{i\bar{j}} \partial_\mu \varphi^i \partial^\mu \varphi^{*\bar{j}}. \end{aligned} \quad (4.3)$$

Although we consider only the scalar part for simplicity, the other parts are consistent with it. We rescale the scalar fields as follows:

$$\frac{1}{\lambda} \varphi = \tilde{\varphi}. \quad (4.4)$$

Then, the Lagrangian (4.3) have the normalized kinetic term:

$$\mathcal{L}_{scalar} = \tilde{g}_{i\bar{j}}(\lambda \tilde{\varphi}, \lambda \tilde{\varphi}^*) \partial_\mu \tilde{\varphi}^i \partial^\mu \tilde{\varphi}^{*\bar{j}}. \quad (4.5)$$

Here,

$$\tilde{g}_{i\bar{j}}|_{\tilde{\varphi}, \tilde{\varphi}^*=0} = \delta_{i\bar{j}} \quad (4.6)$$

is the metric of the manifold of unit radius. Rescaling the WRG eq.(3.31) and comparing the coefficient of $\partial_\mu \tilde{\varphi}^i \partial^\mu \tilde{\varphi}^{*\bar{j}}$, we have

$$\frac{\partial}{\partial t} \tilde{g}_{i\bar{j}}(\lambda \tilde{\varphi}, \lambda \tilde{\varphi}^*) = -\frac{1}{2\pi} \tilde{R}_{i\bar{j}} - \gamma [\tilde{\varphi}^k \tilde{g}_{i\bar{j},k} + \tilde{\varphi}^{*\bar{k}} \tilde{g}_{i\bar{j},\bar{k}} + 2\tilde{g}_{i\bar{j}}],$$

where $\tilde{R}_{i\bar{j}}$ is rescaled Ricci tensor and can be written as

$$\tilde{R}_{i\bar{j}} = h\lambda^2 \tilde{g}_{i\bar{j}} \quad (4.7)$$

using Einstein Kähler condition (4.1).

Because only λ depends on t , this differential equation can be rewritten:

$$\begin{aligned} \frac{\dot{\lambda}}{\lambda} \tilde{\varphi}^k \tilde{g}_{i\bar{j},k} + \frac{\dot{\lambda}}{\lambda} \tilde{\varphi}^{*\bar{k}} \tilde{g}_{i\bar{j},\bar{k}} &= -\left(\frac{h\lambda^2}{2\pi} + 2\gamma\right) \tilde{g}_{i\bar{j}} - \gamma [\tilde{\varphi}^k \tilde{g}_{i\bar{j},k} + \tilde{\varphi}^{*\bar{k}} \tilde{g}_{i\bar{j},\bar{k}}]. \end{aligned} \quad (4.8)$$

Since the left-hand side vanishes for $\varphi, \varphi^* \approx 0$, the coefficient of $\tilde{g}_{i\bar{j}}$ must vanish on the right-hand side. Thus, we obtain the anomalous dimension of scalar fields (or chiral superfields):

$$\gamma = -\frac{h\lambda^2}{4\pi}. \quad (4.9)$$

Comparing the coefficient of $\tilde{\varphi}^k \tilde{g}_{i\bar{j},k}$ (or $\tilde{\varphi}^{*\bar{k}} \tilde{g}_{i\bar{j},\bar{k}}$), we also obtain the β function of λ :

$$\begin{aligned} \beta(\lambda) &\equiv -\frac{d\lambda}{dt} \\ &= -\frac{h}{4\pi}\lambda^3. \end{aligned} \quad (4.10)$$

This β function shows this theory is asymptotically free. The effective coupling constant λ becomes small at the high energy. Thus at UV region the radius $a \rightarrow \infty$ and the manifolds become flat.

The Gell-Mann-Low function based on perturbation theory is discussed in ref.[3]. In that paper, the β functions are calculated using specific forms of Kähler potential for the four group of compact homogeneous symmetric Kähler manifolds. Our method does not need such potential. We found the theories with Einstein Kähler manifolds are asymptotically free in general. To conclude this section, we give two simple examples of Einstein Kähler manifolds. The constant h for the hermitian symmetric space is given by the eigenvalue of the quadratic Casimir operator in adjoint representation of G as is shown in Table 1.

G/H	Dimensions (complex)	h
$SU(N)/[SU(N-1) \otimes U(1)] = CP^{N-1}$	N-1	N
$SU(N)/[SU(N-M) \otimes U(M)]$	M(N-M)	N
$SO(N)/[SO(N-2) \otimes U(1)] = Q^{N-2}$	N-2	N-2
$Sp(N)/U(N)$	$\frac{1}{2}N(N+1)$	N+1
$SO(2N)/U(N)$	$\frac{1}{2}N(N+1)$	N-1
$E_6/[SO(10) \otimes U(1)]$	16	12
$E_7/[E_6 \otimes U(1)]$	27	18

Table 1: The values of h for hermitian symmetric spaces

- CP^N model

The Kähler potential involves a set of N chiral and anti-chiral super-field $\vec{\Phi} = (\Phi^1, \dots, \Phi^N)$

$$K[\Phi, \Phi^\dagger] = a^2 \ln(1 + \vec{\Phi} \vec{\Phi}^\dagger). \quad (4.11)$$

From this Kähler potential, we can obtain Kähler metric and Ricci tensor.

$$\begin{aligned} g_{i\bar{j}} &\equiv \partial_i \partial_{\bar{j}} K \\ &= a^2 \left(\frac{\delta_{i\bar{j}}}{1 + \vec{\varphi} \vec{\varphi}^*} \right) \end{aligned} \quad (4.12)$$

$$\begin{aligned} R_{i\bar{j}} &\equiv -\partial_{\bar{j}} \partial_i (\ln \det g_{k\bar{l}}) \\ &= \frac{N+1}{a^2} g_{i\bar{j}}. \end{aligned} \quad (4.13)$$

Eq.(4.13) shows that this manifold is Einstein Kähler manifold with $h = N + 1$. Hence Eqs.(4.9) and (4.10) give us immediately

$$\gamma = -\frac{(N+1)\lambda^2}{4\pi} \quad (4.14)$$

$$\beta(\lambda) = -\frac{(N+1)\lambda^3}{4\pi}. \quad (4.15)$$

These results are consistent with the perturbative results [3].

- Q^N model

The next example is $Q^N = SO(N+2)/[SO(N) \times U(1)]$ model, whose Kähler potential has the form

$$K[\Phi, \Phi^\dagger] = a^2 \ln \left(1 + \vec{\Phi} \vec{\Phi}^\dagger + \frac{1}{4} \vec{\Phi}^2 \vec{\Phi}^{\dagger 2} \right), \quad (4.16)$$

where $\vec{\Phi} = (\Phi^1, \dots, \Phi^N)$. Kähler metric and Ricci tensor are given by

$$\begin{aligned} g_{i\bar{j}} &= a^2 \left[\frac{\delta_{i\bar{j}}}{1 + \vec{\varphi} \vec{\varphi}^* + \frac{1}{4} \vec{\varphi}^2 \vec{\varphi}^{*2}} \right. \\ &\quad \left. + \frac{\varphi^i \varphi^{*\bar{j}} (1 + \vec{\varphi} \vec{\varphi}^*) - \left(\varphi_i^* \varphi_{\bar{j}} + \frac{1}{2} \vec{\varphi}^2 \varphi_i^* \varphi^{*\bar{j}} + \frac{1}{2} \vec{\varphi}^{*2} \varphi^i \varphi_{\bar{j}} \right)}{\left(1 + \vec{\varphi} \vec{\varphi}^* + \frac{1}{4} \vec{\varphi}^2 \vec{\varphi}^{*2} \right)^2} \right], \end{aligned} \quad (4.17)$$

$$R_{i\bar{j}} = \frac{N}{a^2} g_{i\bar{j}}. \quad (4.18)$$

Eq.(4.18) shows that this manifold is also Einstein Kähler manifold, and $h = N$. According to the same discussion as CP^N model, we obtain the anomalous dimension and β function:

$$\gamma = -\frac{N\lambda^2}{4\pi} \quad (4.19)$$

$$\beta(\lambda) = -\frac{N\lambda^3}{4\pi}. \quad (4.20)$$

Combining these two examples, we will construct a new model in the next section.

5 A New Model

Consider the following Kähler potential which has two parameters $\lambda(t) = \frac{1}{a(t)}$ and $g(t)$:

$$K[\Phi, \Phi^\dagger] = a^2(t) \ln \left(1 + \vec{\Phi} \vec{\Phi}^\dagger + g(t) \vec{\Phi}^2 \vec{\Phi}^{\dagger 2} \right). \quad (5.1)$$

When $g = 0$, this is CP^N model, and this becomes Q^N model when $g = \frac{1}{4}$. The former has the global symmetry $SU(N+1)$, and the latter has $SO(N+2)$.

This Kähler potential gives the Kähler metric and Ricci tensor:

$$\begin{aligned} g_{i\bar{j}} &= a^2(t) \left[\frac{\delta_{i\bar{j}}}{1 + \vec{\varphi} \vec{\varphi}^* + g(t) \vec{\varphi}^2 \vec{\varphi}^{*2}} \right. \\ &\quad \left. + \frac{4g(t) \varphi^i \varphi^{*\bar{j}} (1 + \vec{\varphi} \vec{\varphi}^*) - (\varphi_i^* \varphi_{\bar{j}} + 2g(t) \vec{\varphi}^2 \varphi_i^* \varphi^{*\bar{j}} + 2g(t) \vec{\varphi}^{*2} \varphi^i \varphi_{\bar{j}})}{(1 + \vec{\varphi} \vec{\varphi}^* + g(t) \vec{\varphi}^2 \vec{\varphi}^{*2})^2} \right], \\ R_{i\bar{j}} &= (N+1) \frac{1}{a(t)} g_{i\bar{j}} - \left[\frac{4g(t) \delta_{i\bar{j}}}{1 + 4g(t) \vec{\varphi} \vec{\varphi}^* + g(t) \vec{\varphi}^2 \vec{\varphi}^{*2}} \right. \\ &\quad \left. + \frac{16g^2(t) \varphi^i \varphi^{*\bar{j}} \vec{\varphi} \vec{\varphi}^* - 16g^2(t) \varphi_i^* \varphi_{\bar{j}} - 8g^2(t) (\vec{\varphi}^2 \varphi_i^* \varphi^{*\bar{j}} + \vec{\varphi}^{*2} \varphi^i \varphi_{\bar{j}})}{(1 + 4g(t) \vec{\varphi} \vec{\varphi}^* + g(t) \vec{\varphi}^2 \vec{\varphi}^{*2})^2} \right]. \end{aligned} \quad (5.2)$$

Note that Eq.(5.2) shows this manifold is not an Einstein Kähler manifold unless g takes specific values. According to the same discussion as section 4, we obtain

$$\gamma = -\frac{\lambda^2}{4\pi} [(N+1) - 4g], \quad (5.3)$$

$$\beta(\lambda) = -\frac{\lambda^3}{4\pi} [(N+1) + 8g(2g-1)], \quad (5.4)$$

$$\beta(g) = \frac{4\lambda^2}{\pi} g^2 (4g-1). \quad (5.5)$$

Fig.3 shows the renormalization group flows in $\lambda - g$ plane.

From Eq.(5.5), $\beta(g)$ is zero when $g = 0$ and $g = \frac{1}{4}$. These values of g correspond to CP^N and Q^N models respectively, and in both cases the manifolds become Einstein Kähler manifolds.

In the infrared(IR) region, the coupling constant λ becomes infinity and g approaches $\frac{1}{4}$. Thus this model becomes Q^N model with strong coupling at low energy. On the contrary, at ultraviolet(UV) region, the flow separates $g < \frac{1}{4}$ and $g > \frac{1}{4}$ regions. In $g < \frac{1}{4}$ region, the coupling constant $\lambda \rightarrow 0$ and $g \rightarrow 0$. Thus CP^N with the radius $a \rightarrow \infty$ is the UV fixed point of this new model. In $g > \frac{1}{4}$ region, the coupling constant g goes to infinity and λ becomes zero.




Figure 3: Renormalization group flows (The direction of arrows shows infrared region.)

6 Conclusions

In this paper we have found the β function for the WRG equation in two dimensional $\mathcal{N} = 2$ supersymmetric nonlinear sigma model using nonperturbative method. This β function of Kähler metric at $O(\partial^2)$ approximation consists of the one-loop correction term and rescaling term. We have obtained several new results from this β function.

First, we have seen all theories are asymptotically free if target spaces are generic compact Einstein Kähler manifolds. This statement is drawn by using only Einstein Kähler condition, and does not depend on the specific form of Kähler potential.

Second, we have constructed a new model, which has two parameters g and λ . This model is not Einstein Kähler manifold except for special values of g . Using the Wilsonian renormalization group equation, we obtain two β functions of each parameters. One of the β functions ($\beta(g)$) is zero when the model becomes CP^N and Q^N model. These two models have the different global symmetry $SU(N+1)$ and $SO(N+2)$ respectively, and both of them are Einstein Kähler manifolds. The renormalization group flow shows the new model interpolates CP^N model and Q^N model. We have also found the CP^N model is the UV fixed point of the model. From this analysis, we conjecture that generic Kähler manifolds go to Einstein Kähler manifolds through Wilsonian renormalization group flow.

Acknowledgements

We would like to thank Muneto Nitta for useful communication in early stage of this work.

References

- [1] A. D’Adda, P. Di Vecchia and M. Luscher, Confinement And Chiral Symmetry Breaking In CP^{N-1} Models With Quarks Nucl. Phys. B **152** (1979) 125.
K.Higashijima, T.Kimura, M.Nitta and M.Tsuzuki, Large-N limit of $N = 2$ supersymmetric Q^N model in two dimensions Prog. Theor. Phys. **105** (2001) 261 [arXiv:hep-th/0010272]
- [2] K.Higashijima, T.Kimura and M.Nitta, Gauge theoretical construction of non-compact Calabi-Yau manifolds To appear in Ann.of Phys [arXiv:hep-th/0110216].
K.Higashijima, T.Kimura and M.Nitta, Calabi-Yau Manifolds of Cohomogeneity One as Complex Line Bundles [arXiv:hep-th/0202064].
- [3] A.Y.Morozov, A.M.Perelomov and M.A.Shifman, Exact Gell-Mann-Low Function Of Supersymmetric Kahler Sigma Models Nucl. Phys. B **248** (1984) 279
- [4] K.G.Wilson and I.G.Kogut, The Renormalization Group And The ϵ Expansion Phys.Rep.**12**,75 (1974)
- [5] F.Wegner and A.Houghton, Renormalization Group equation for Critical Phenomena Phys.Rev.**A8**,401(1973)
- [6] T.R.Morris, The Exact renormalization group and approximate solutions Int. J. Mod. Phys. A **9** (1994) 2411 [arXiv:hep-ph/9308265].
T.R.Morris, Derivative expansion of the exact renormalization group Phys. Lett. B **329** (1994) 241 [arXiv:hep-ph/9403340].
T.R.Morris and M.D.Turner, Derivative expansion of the renormalization group in $O(N)$ scalar field theory Nucl. Phys. B **509** (1998) 637 [arXiv:hep-th/9704202].
- [7] K.Aoki, Introduction To The Non-Perturbative Renormalization Group And Its Recent Applications Int. J. Mod. Phys. B **14** (2000) 1249.
- [8] T.E.Clark, B.Haeri and S.T.Love, Wilson Renormalization Group Analysis Of Theories With Scalars And Fermions Nucl. Phys. B **402** (1993) 628 [arXiv:hep-ph/9211261].

- [9] T.E.Clark and S.T.Love, The Wilson effective Kaehler potential for supersymmetric nonlinear sigma models Phys. Rev. D **56** (1997) 2461 [arXiv:hep-th/9701134].
- [10] K.Higashijima and M.Nitta, Kaehler normal coordinate expansion in supersymmetric theories Prog. Theor. Phys. **105** (2001) 243 [arXiv:hep-th/0006027]
- [11] K.Higashijima,E.Itou and M.Nitta
Normal Coordinates in Kähler Manifolds and the Background Field Method hep-th/0203081
- [12] L.Alvarez-Gaumé,D.Z.Freedman and S.Mukhi, The Background Field Method And The Ultraviolet Structure Of The Supersymmetric Non-linear σ -Model Ann.of Phys. **134** (1981) 85
- [13] D.G.Boulware and L.S.Brown, Symmetric Space Scalar Field Theory Ann.of Phys. **138** (1982) 392